

Robustness of the Bennett-Brassard 1984 quantum key distribution protocol against general coherent attacks

Georgios M. Nikolopoulos and Gernot Alber

Institut für Angewandte Physik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

(Dated: February 1, 2008)

It is demonstrated that for the entanglement-based version of the Bennett-Brassard (BB84) quantum key distribution protocol, Alice and Bob share provable entanglement if and only if the estimated qubit error rate is below 25% or above 75%. In view of the intimate relation between entanglement and security, this result sheds also new light on the unconditional security of the BB84 protocol in its original prepare-and-measure form. In particular, it indicates that for small qubit error rates 25% is the ultimate upper security bound for any prepare-and-measure BB84-type QKD protocol. On the contrary, for qubit error rates between 25% and 75% we demonstrate that the correlations shared between Alice and Bob can always be explained by separable states and thus, no secret key can be distilled in this regime.

PACS numbers: 03.67.Dd, 03.65.Ud

I. INTRODUCTION

Quantum cryptography solves an outstanding problem of classical cryptography, namely the distribution of a secret perfectly correlated random key between legitimate users. The first quantum key distribution (QKD) protocol, which was able to achieve this goal by exploiting characteristic quantum phenomena, was proposed by Bennet and Brassard [1]. This BB84 protocol is the prototype of a prepare-and-measure QKD protocol, which allows unconditionally secure key distribution between two legitimate users, Alice and Bob. Its security against an eavesdropper, Eve, is guaranteed by the peculiar features of the quantum mechanical measurement process [2]. Later, it was realized that the security of the BB84 protocol is based on an effective entanglement shared between Alice and Bob. In particular, it was shown that, from the point of view of Eve, the originally proposed prepare-and-measure scheme is equivalent to an associated entanglement-based version of the BB84 protocol [3, 4, 5, 6]. The security of this latter protocol is guaranteed, for example, if the state of the entangled-qubit pairs shared between Alice and Bob is exponentially close to being a pure state. Then, according to Holevo's theorem, Eve's mutual information with the distributed key is exponentially small [4, 6].

This equivalence between a prepare-and-measure QKD protocol and its associated entanglement-based QKD protocol, has been exploited in various proofs of unconditional security of the BB84 protocol [4, 5, 6]. Indeed, that quantum entanglement is necessary for secret QKD, was also emphasized in a more general context, recently [7, 8]. Nevertheless, though with respect to unconditional security issues the prepare-and-measure and the entanglement-based forms of the BB84 protocol are equivalent, as far as realistic implementations are concerned both schemes are quite different. Specifically, entanglement-based QKD protocols typically require quantum memories and quantum computers for performing the entanglement purification, necessary for

enforcing secrecy even in the presence of an arbitrary powerful eavesdropper. On the contrary, the corresponding prepare-and-measure schemes do not require this advanced quantum technology, and are therefore realizable much more easily with present day technology.

An important quantity characterizing the security of a QKD protocol is the *threshold disturbance*, i.e. the maximal disturbance or quantum bit error rate (QBER) which can be tolerated by Alice and Bob for being capable of producing a secret, perfectly correlated random key. This maximal disturbance quantifies the robustness of a QKD scheme, and in general it depends not only on the eavesdropping strategy, but also on the algorithm (distillation protocol) Alice and Bob are using for post-processing their raw data [9]. From the practical point of view, it is most interesting to know the threshold disturbance of a given QKD protocol, with respect to the most general eavesdropping attacks which are consistent with the laws of quantum mechanics. For such general coherent attacks, it has been demonstrated, for example, that the prepare-and-measure scheme of the BB84 protocol is unconditionally secure up to QBERs of approximately 20%, provided one uses appropriate error correction and privacy amplification protocols, which involve two-way classical communication [10, 11]. Furthermore, it is also known that a simple intercept-resend attack can beat the BB84 protocol giving rise to QBERs $\geq 25\%$ [12]. However, so far it is unknown whether an arbitrary powerful eavesdropper, whose power is only limited by the general laws of quantum mechanics, is able to push the threshold disturbance below 25% [9, 11, 13].

In this paper we investigate the robustness of the BB84 protocol, with respect to general coherent attacks by an adversary whose power is only limited by the fundamental laws of quantum mechanics. As a main result, it is shown that for QBERs less than 25% Alice and Bob always share pairs of entangled qubits, even if Eve has performed a general coherent attack involving quantum memories and quantum computers. In other words, as far as the robustness of the protocol is concerned, the

coherent attacks do not seem to offer any advantage over collective or incoherent attacks. For the entanglement-based version of the BB84 protocol our result implies that Alice and Bob can distribute a random secret key up to this QBER, provided they purify their corrupted pairs with the help of an entanglement purification protocol (EPP) [14, 15]. This result also sheds new light onto the still open question of the maximal error rate that can be tolerated by a prepare-and-measure scheme of the BB84 protocol [11, 13]. In particular, in view of the equivalence between the prepare-and-measure scheme and the entanglement-based version of the BB84 protocol, our result demonstrates that 25% is indeed the threshold disturbance for the prepare-and-measure scheme as well. Thus, in principle Alice and Bob are able to exploit these quantum correlations for the distillation of a perfectly correlated secret random key, by means of an appropriate classical distillation protocol which involves two-way classical communication [10, 11]. Furthermore, as long as Alice and Bob focus on their sifted key and manipulate each qubit-pair independently, for QBERs between 25% and 75% a secret-key distillation is impossible. In particular, in this case we show that the correlations shared between Alice and Bob can always be described by separable states and thus the necessary precondition for secret-key distillation is violated [7, 8].

This paper is organized as follows: For the sake of completeness in Sec. II we summarize briefly basic facts of the prepare-and-measure scheme and the equivalent entanglement-based version of the BB84 protocol, as far as they are significant for our subsequent discussion. In Sec. III we present the proof of our main result while Sec. IV contains discussion and conclusions.

II. BASIC FACTS ABOUT BB84 PROTOCOL

In the prepare-and-measure BB84 protocol, Alice sends a sequence of qubits to Bob each of which is randomly prepared in one of the basis states $\{|0\rangle, |1\rangle\}$ or $\{|\bar{0}\rangle, |\bar{1}\rangle\}$ which are eigenstates of two maximally conjugate physical variables, namely the two Pauli spin operators $\mathcal{Z}$ and $\mathcal{X}$. The eigenstates of $\mathcal{Z}$, i.e. $\{|0\rangle, |1\rangle\}$, and of $\mathcal{X}$, i.e. $\{|\bar{0}\rangle, |\bar{1}\rangle\}$, are related by the Hadamard transformation

$$\mathcal{H} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad (1)$$

i.e. $|\bar{i}\rangle = \sum_j \mathcal{H}_{ij} |j\rangle$ ($i, j = 0, 1$). In the basis $\{|0\rangle, |1\rangle\}$, the Pauli spin operators are represented by the matrices

$$\mathcal{X} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \mathcal{Y} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \mathcal{Z} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2)$$

Bob measures the received qubits randomly in one of the two bases. Afterwards Alice and Bob apply a random permutation of their data, and publicly discuss the bases chosen, discarding all the bits where they have selected

different bases. Subsequently, they randomly select half of the bits from the remaining random key (sifted key) and determine its error probability or QBER. If, as a result of a noisy quantum channel or of an eavesdropper, the estimated QBER is too high, they abort the protocol. Otherwise, they perform error correction and privacy amplification with one- or two-way classical communication, in order to obtain a smaller number of secret and perfectly correlated random bits.

It has been shown that, from the point of view of an arbitrarily powerful eavesdropper, this originally proposed prepare-and-measure form of the BB84 protocol is equivalent to an entanglement-based QKD protocol [3, 4, 5, 6]. This latter form of the BB84 protocol offers advantages in particular with respect to questions concerning its unconditional security, and works as follows: Alice prepares each of, say $2n$, entangled-qubit pairs in a particular Bell state [16], say $|\Phi^+\rangle \equiv \frac{1}{\sqrt{2}}(|0_A 0_B\rangle + |1_A 1_B\rangle)$ (where the subscripts A, B refer to Alice and Bob, respectively). After having applied Hadamard transformation on the second half of each pair of a subset of randomly chosen qubit pairs, she sends all the second halves of the $2n$ pairs to Bob. Due to channel noise and the presence of a possible eavesdropper, however, at the end of this transmission stage all the $2n$ entangled-qubit pairs will be corrupted. In fact, they will be entangled among themselves as well as with Eve's probe.

In order to ensure secret key distribution even in the presence of such a powerful eavesdropper, first of all Alice and Bob perform a *random permutation* of all the pairs, thus distributing any influence of the channel noise or the eavesdropper, equally among all the pairs [5, 11]. Afterwards, they perform a verification test, in which they randomly select half of the pairs [17] as check pairs and measure each one of them *independently*, along a common basis, i.e. the quantization axis of the Bell state $|\Phi^+\rangle$. However, these measurements are done only after Bob has reversed Alice's randomly applied Hadamard transformations. For this latter purpose, after the transmission stage of the protocol, Alice has to announce publicly the qubits on which she has applied Hadamard transformations. Subsequently, the influence of channel noise or of an eavesdropper is quantified by the average QBER of these n check pairs, i.e. the probability with which Alice's and Bob's measurement results disagree. Note that, since Alice and Bob perform their measurements along the quantization axis of the Bell state $|\Phi^+\rangle$, in the absence of noise and eavesdropping their measurement results would perfectly agree. Furthermore, assuming that the check pairs constitute a fair sample [17], one may conclude that the same QBER applies also to the remaining, yet unmeasured, n pairs. After this verification test all the check pairs are dismissed and, if the QBER is too high, the protocol is aborted. Otherwise, Alice and Bob proceed to the purification of the remaining n pairs

For this purpose, they apply an appropriate entanglement purification protocol (EPP) [14, 15] with classical one- or two-way communication, so that they are able

to distill a smaller number of almost pure entangled-qubit pairs. Finally, measuring these almost perfectly entangled-qubit pairs in a common basis, i.e. along the quantization axis of the Bell state $|\Phi^+\rangle$, Alice and Bob obtain a perfectly correlated random key, about which an adversary has negligible information. For the sake of completeness we finally want to mention that this entanglement-based protocol can be reduced to an associated prepare-and-measure BB84 protocol with error correction and privacy amplification involving one- or two-way classical communication, only if the underlying EPP fulfills certain consistency conditions which are of no concern for our further discussion [5, 10, 11].

On the basis of the equivalence of these two forms of the BB84 protocol, it has been demonstrated that by means of one-way classical distillation protocols a random key can be exchanged secretly between the two legitimate users up to a QBER of approximately 11% [5]. Furthermore, post-processing that involves two-way classical communication can guarantee the extraction of a secret key even up to QBERs of the order of 20% [10, 11]. On the other hand, it is also known that an intercept-resend attack can always defeat the BB84 protocol (irrespective of the distillation protocol employed by Alice and Bob) for QBERs above 25% [12]. However, as far as QBERs between 20% and 25% are concerned, it is still an open question whether Alice and Bob are able to exchange a secret key, under the assumption of general coherent attacks.

III. ENTANGLEMENT AND THRESHOLD DISTURBANCE OF THE BB84 PROTOCOL

According to a recent observation, a *necessary precondition* for secret key distillation is that the correlations

established between Alice and Bob during the state distribution cannot be explained by a separable state [7, 8]. Throughout this work, we consider that Alice and Bob focus on the sifted key during the post processing (i.e., they discard immediately all the polarization data for which they have used different bases) and that they treat each pair independently. Thus, according to the aforementioned precondition, given a particular value of the estimated QBER, the task of Alice and Bob is to infer whether they share provable entanglement or not. Thereby, entanglement is considered to be provable if Alice's and Bob's correlations cannot be explained by a separable state within the framework of the protocol considered. In this section we demonstrate for the BB84 protocol that, as long as the detected QBER is less than 25% or larger than 75%, Alice and Bob can be confident that they share entangled-qubit pairs with high probability. Furthermore, for QBERs between 25% and 75% the correlations between Alice and Bob can always be explained by a separable two-qubit state.

Adopting the entanglement-based version of the BB84 protocol we described in the previous section, the average disturbance that Alice and Bob estimate during the verification test is given by [5, 11, 19]

$$\text{QBER} = \frac{1}{2} \sum_{b=0,1} \sum_{l=0,1} \frac{1}{n} \sum_{j_i; i=1}^n \text{Tr}_{A,B} \{ [\mathcal{H}_A^b | l \rangle_{AA} \langle l | \mathcal{H}_A^b \otimes \mathcal{H}_B^b | l \oplus 1 \rangle_{BB} \langle l \oplus 1 | \mathcal{H}_B^b]_{j_i} \rho_{AB} \}, \quad (3)$$

where $\oplus$ denotes addition modulo 2. Thereby ρ_{AB} denotes the reduced density operator of Alice and Bob for all $2n$ pairs while the index j_i indicates that the corresponding physical observable refers to the j_i -th randomly selected qubit pair. The powers of the Hadamard transformation $\mathcal{H}^b$, with $b \in \{0, 1\}$, reflect the fact that the errors in the sifted key originate from measurements in both complementary bases which have been selected randomly by Alice and Bob with equal probabilities.

Recall now that Alice and Bob have permuted all their pairs randomly before the verification stage. This random permutation has the effect of distributing the errors introduced by Eve, over all the qubit pairs in a homo-

geneous way [5, 11]. As a result of this *homogenization procedure* all the reduced density operators describing the state of each pair shared between Alice and Bob become equal, i.e.

$$\rho_{AB}^{(1)} = \rho_{AB}^{(2)} = \dots = \rho_{AB}^{(2n)}. \quad (4)$$

The reduced density operator of Alice's and Bob's k -th pair is denoted by $\rho_{AB}^{(k)} = \text{Tr}_{AB}^{(\neq)}(\rho_{AB})$ and $\text{Tr}_{AB}^{(\neq)}$ indicates the tracing (averaging) procedure over all the qubit pairs except the k -th one. Note that in general ρ_{AB} is expected to have a complicated structure as it includes all the effects arising from a general coherent attack of a

possible eavesdropper. As a result of the homogenization procedure, however, the reduced states of the pairs $\rho_{AB}^{(k)}$

are equal and thus the average estimated disturbance (3) reads

$$\text{QBER} = \frac{1}{2} \sum_{b=0,1} \sum_{l=0,1} \text{Tr}_{A,B}^{(j_1)} \{ [\mathcal{H}_A^b | l \rangle_{AA} \langle l | \mathcal{H}_A^b \otimes \mathcal{H}_B^b | l \oplus 1 \rangle_{BB} \langle l \oplus 1 | \mathcal{H}_B^b]_{j_1} \rho_{AB}^{(j_1)} \}, \quad (5)$$

where $\text{Tr}_{A,B}^{(j_1)}$ denotes the tracing procedure over the j_1 -th qubit pair of Alice and Bob. According to Eq. (5), and in view of the homogenization procedure, the QBER is determined by the *average error probability* of an arbitrary qubit pair, say the pair j_1 . Thus, the corresponding reduced density operator $\rho_{AB}^{(j_1)}$ contains all the necessary information about the noisy quantum channel and about a possible general coherent attack by an eavesdropper, which is relevant for the evaluation of the average QBER. In particular, this implies that an arbitrary attack by an eavesdropper which gives rise to a particular state $\rho_{AB}^{(j_1)}$ is indistinguishable, from the point of view of the estimated average disturbance, from a corresponding individual attack which results in a decorrelated n -pair state of the form $\bigotimes_{i=1}^n \rho_{AB}^{(j_i)}$.

In order to address the problem for which values of the QBER the reduced density operator $\rho_{AB}^{(j_1)}$ is entangled, we start from the observation that the QBER of Eq. (5) is invariant under the transformations

$$\begin{aligned} (l, b) &\rightarrow (l \oplus 1, b), \\ (l, b) &\rightarrow (l, b \oplus 1). \end{aligned} \quad (6)$$

This invariance implies that there are various reduced density operators of the j_1 -th qubit pair, which all give rise to the same observed value of the QBER. This can be seen from the elementary relations

$$\begin{aligned} \mathcal{X} \mathcal{H}^b | l \rangle \langle l | \mathcal{H}^b \mathcal{X} &= \mathcal{H}^b | l \oplus 1 \oplus b \rangle \langle l \oplus 1 \oplus b | \mathcal{H}^b, \\ \mathcal{Z} \mathcal{H}^b | l \rangle \langle l | \mathcal{H}^b \mathcal{Z} &= \mathcal{H}^b | l \oplus b \rangle \langle l \oplus b | \mathcal{H}^b, \\ \mathcal{X} \mathcal{Z} \mathcal{H}^b | l \rangle \langle l | \mathcal{H}^b \mathcal{Z} \mathcal{X} &= \mathcal{H}^b | l \oplus 1 \rangle \langle l \oplus 1 | \mathcal{H}^b. \end{aligned} \quad (7)$$

Together with the invariance of the QBER under the transformations (6), these relations imply that the reduced operators $\rho_{AB}^{(j_1)}$ and

$$\tilde{\rho}_{AB}^{(j_1)} = \frac{1}{8} \sum_{g \in \mathcal{G}_1, h \in \mathcal{G}_2} U(h) U(g) \rho_{AB}^{(j_1)} U(g)^\dagger U(h)^\dagger \quad (8)$$

give rise to the same value of the QBER. Thereby, the unitary and hermitian operators

$$\begin{aligned} U(g_1) &= \mathcal{X}_A \otimes \mathcal{X}_B, \\ U(g_2) &= \mathcal{Z}_A \otimes \mathcal{Z}_B, \\ U(g_3) &= -\mathcal{Y}_A \otimes \mathcal{Y}_B \equiv \mathcal{X}_A \mathcal{Z}_A \otimes \mathcal{X}_B \mathcal{Z}_B, \\ U(g_4) &= \mathbb{1}_A \otimes \mathbb{1}_B, \end{aligned} \quad (9)$$

and

$$\begin{aligned} U(h_1) &= \mathcal{H}_A \otimes \mathcal{H}_B, \\ U(h_2) &= \mathbb{1}_A \otimes \mathbb{1}_B, \end{aligned} \quad (10)$$

have been introduced, which form unitary representations of two discrete Abelian groups $\mathcal{G}_1 = \{g_1, g_2, g_3, g_4\}$ and $\mathcal{G}_2 = \{h_1, h_2\}$. Invariance of $\tilde{\rho}_{AB}^{(j_1)}$ under these groups is induced by the symmetry transformations of Eq. (6) which leave the QBER invariant.

As a consequence of Eq. (8), the density operator $\rho_{AB}^{(j_1)}$ is entangled if $\tilde{\rho}_{AB}^{(j_1)}$ is entangled, as both states are related by local unitary operations and convex summation. Hence, to determine the values of the QBER for which Alice and Bob share provable entanglement, it suffices to determine the QBERs for which the most general two-qubit state $\tilde{\rho}_{AB}^{(j_1)}$ (which is invariant under the Abelian discrete groups $\mathcal{G}_1$ and $\mathcal{G}_2$) is entangled.

Note now that, the hermitian operators $U(g_1)$ and $U(g_2)$ of the group $\mathcal{G}_1$ constitute already a complete set of commuting operators in the Hilbert space of two qubits. Their eigenstates are given by the Bell states [16]. Thus, the most general two-qubit state which is invariant under the Abelian group $\mathcal{G}_1$ is given by a convex sum of Bell states, i.e.

$$\tilde{\rho}_{AB}^{(j_1)} = \lambda_{00} |\Phi^+\rangle \langle \Phi^+| + \lambda_{10} |\Phi^-\rangle \langle \Phi^-| + \lambda_{01} |\Psi^+\rangle \langle \Psi^+| + \lambda_{11} |\Psi^-\rangle \langle \Psi^-|. \quad (11)$$

Thereby, the non-negative parameters $\lambda_{\alpha\beta}$ have to fulfill

the normalization condition

$$\sum_{\alpha, \beta \in \{0,1\}} \lambda_{\alpha\beta} = 1. \quad (12)$$

Furthermore, additional invariance of the quantum state (11) under the discrete group $\mathcal{G}_2$ implies that

$$\lambda_{01} = \lambda_{10}. \quad (13)$$

As a consequence of this last constraint and the normalization condition, only two of the four parameters $\lambda_{\alpha\beta}$ are independent. According to Eq.(5), the average QBER is related to these parameters by

$$\text{QBER} = \lambda_{11} + \lambda_{01}. \quad (14)$$

Let us now determine the possible values of the QBER for which Alice and Bob share provable entanglement. To this end, we can take advantage of the Peres-Horodecki criterion, for example, stating that the state $\tilde{\rho}_{AB}^{(j_1)}$ is disentangled *iff* its partial transpose is non-negative [20]. In our case, this condition is equivalent to the inequalities

$$\lambda_{01} + \lambda_{11} \geq |\lambda_{00} - \lambda_{10}|, \quad (15)$$

$$\lambda_{00} + \lambda_{10} \geq |\lambda_{01} - \lambda_{11}|, \quad (16)$$

As depicted in Fig. 1, these inequalities, combined with Eqs. (12) and (13), imply that the reduced den-

sity operator $\tilde{\rho}_{AB}^{(j_1)}$ is entangled if the estimated QBER is below 1/4 or above 3/4. Given, however, that the states $\tilde{\rho}_{AB}^{(j_1)}$ and $\rho_{AB}^{(j_1)}$ are related via local operations and convex summation, the original single-pair state $\rho_{AB}^{(j_1)}$ must also be entangled in the same regime of parameters. Hence we may conclude that, whenever Alice and Bob detect an average QBER within this interval, they can be confident that they share entanglement with high probability, and their correlations cannot originate from a separable state. The necessary precondition for secret-key distillation is therefore fulfilled.

Let us now consider average QBERs with $1/4 \leq \text{QBER} \leq 3/4$. The fact that $\tilde{\rho}_{AB}^{(j_1)}$ is separable does not necessarily imply that $\rho_{AB}^{(j_1)}$ is also separable. But it indicates that in this regime of parameters, Alice's and Bob's correlations within the framework of the BB84 protocol can be explained by a separable state, namely by $\tilde{\rho}_{AB}^{(j_1)}$. According to [7, 8], this implies that Alice and Bob cannot extract a secret key. A particularly simple family of such separable states defined for $1/4 \leq \text{QBERs} \leq 3/4$, for example, is given by

$$\sigma_{AB}(D) = (4D - 1)\frac{\mathbb{1}_4}{4} + |1 - 2D| \sum_{k \in \{0,1\}} \left[\frac{1}{2} (|k_A\rangle\langle k_A|) \otimes (|k_B\rangle\langle k_B|) + (\tilde{\sigma}_A^{(k)} \otimes \tilde{\sigma}_B^{(k)}) \right]. \quad (17)$$

Thereby, $D \equiv \text{QBER}$, $\mathbb{1}_4$ denotes the unit operator in $\mathbb{C}_A^2 \otimes \mathbb{C}_B^2$ and $\tilde{\sigma}_C^{(k)} = \frac{1}{2} (|0_C\rangle\langle k_C| + |1_C\rangle\langle (1 \oplus k)_C|)$.

The family (17) is parameterized by the estimated average disturbance detected by Alice and Bob. Moreover, any separable state which belongs to this family is indistinguishable, from the point of view of the estimated QBER, from the real state shared between Alice and Bob. In other words, whenever the detected QBER is between 1/4 and 3/4, the correlations shared between Alice and Bob can be very well described in the framework of the family of separable states $\sigma_{AB}(D)$. The necessary precondition [7, 8, 22] for secret key distillation is not met for disturbances within this regime, so that the protocol must be aborted.

IV. DISCUSSION AND CONCLUSIONS

We have discussed the threshold disturbance of the BB84 QKD protocol under the assumption of general coherent attacks which are only limited by the fundamental laws of quantum theory. If Eve disentangles Alice and Bob by such a general attack (i.e., Alice's and Bob's correlations can be explained by a separable state) this gives rise to QBERs between 25% and 75%. Hence, if

Alice and Bob detect a QBER within this interval, they have to abort the protocol. As an example, we presented a particular class of separable states which is capable of describing the correlations shared between Alice and Bob for $25\% \leq \text{QBER} \leq 75\%$. For QBERs below 25% and above 75% the two legitimate users can be confident that with high probability the quantum correlations they share cannot be described by separable states.

The relevance of our results in the context of the entanglement-based version of the BB84 protocol is apparent : Alice and Bob are able to extract a secret key for QBERs $< 25\%$ and QBERs $> 75\%$ by application of an EPP which purifies towards the Bell states $|\Phi^+\rangle$ or $|\Psi^-\rangle$, respectively.

However, for the prepare-and-measure forms of the BB84 protocol there are still open questions. So far, the highest tolerable error rates that have been reported in the context of general coherent eavesdropping attacks are close to 20% [10, 11]. Nevertheless, according to a recent observation [7, 8], a necessary precondition for secret-key distillation is that the correlations established between Alice and Bob during the state distribution cannot be explained by a separable state. In view of this strong link between quantum and secret correlations, our results suggest that in principle the distillation of a secret key




FIG. 1: Regions of the independent parameters $D = \text{QBER}$ and $G = \lambda_{00} - \lambda_{10}$ for which the qubit pair state $\tilde{\rho}_{AB}^{(j_1)}$ is separable (shaded regions). The various constraints that these parameters have to satisfy, are indicated by straight dashed lines, namely (a) Eq. (15); (b) $\lambda_{00}, \lambda_{10} \geq 0$; (c) Eqs. (12), (13) and (16); (d) $\lambda_{01}, \lambda_{11} \geq 0$ and Eqs. (12), (13). The protocol operates in the region which is defined by the solid lines. The threshold disturbances are indicated by black dots while the dot-dashed line connecting these dots corresponds to the family of separable states (17).

is in principle possible for QBERs below 25% and above 75%. In fact, if we focus on low disturbances, we can say that 25% is the upper bound on the maximal QBER that can be tolerated by any BB84-type prepare-and-measure QKD protocol. In other words, as far as the robustness of the BB84 protocol is concerned, coherent attacks do not seem to offer any advantages over collective or even incoherent attacks. Nevertheless, finding error correction and privacy amplification protocols required for a BB84-type protocol to be capable of meeting this upper security bound remains an open problem and will be addressed in subsequent work.

Finally, it should be stressed that the disturbance thresholds we have obtained depend on the post-processing of the BB84 QKD protocol. In particular, they rely on the complete omission of those qubits of the raw key for which Alice and Bob measured in different bases. Furthermore, they also rely on the fact that Alice

and Bob manipulate each qubit pair separately. At first sight it might appear to be puzzling why, under the same assumptions, it was shown that a secret key can be distilled in the framework of the BB84 protocol for QBERs up to about 30%, by means of advantage distillation protocols. These results, however, have been obtained under the additional assumption that Eve is restricted to so-called “optimal incoherent attacks”, i.e. attacks which maximize Eve’s information gain. One can easily verify, for example, that the class of separable states (17) is not optimal (in the sense of [9, 21]). In our work we allow for arbitrary eavesdropping attacks and thus we have demonstrated that the distillation of a secret key for QBERs above 25% is impossible [7, 8, 22]. So, although the incoherent attacks considered in [9, 21] are optimal with respect to the information gain of an eavesdropper [2], they are not able to disentangle Alice and Bob at the lowest possible QBER (that is at 25%). The cost of information loss that Eve has to accept by employing an attack that disentangles Alice and Bob at each particular disturbance between 25% and 75% remains an open question. Clearly, to this end one has to consider in detail the eavesdropping attack and this is beyond the purpose of this work.

Nevertheless, in principle the distillation of a secret key for QBERs above 25% is possible, provided that Alice and Bob do not focus on their sifted key only. In this context it was demonstrated with the help of entanglement witnesses recently, that the detection of quantum correlations between Alice and Bob is feasible even for QBERs above 25%, provided the data of the raw key are also taken into account [7].

V. ACKNOWLEDGMENTS

Stimulating discussions with Norbert Lütkenhaus are gratefully acknowledged. We thank an anonymous referee for insightful comments on the first version of this paper. This work is supported by the EU within the IP SECOQC.

-
- [1] C. H. Bennett and G. Brassard, in *Proceedings of the IEEE International Conference on Computers, Systems and Signal Processing, Bangalore, India* (IEEE, New York, 1984), pp. 175-179.
 - [2] C. A. Fuchs, N. Gisin, R. B. Griffiths, C. S. Niu, and A. Peres, *Phys. Rev. A* **56**, 1163 (1997).
 - [3] C. H. Bennett, G. Brassard, and N. D. Mermin, *Phys. Rev. Lett.* **68**, 557 (1992).
 - [4] H.-K. Lo and H. F. Chau, *Science* **283**, 2050 (1999).
 - [5] P. W. Shor and J. Preskill, *Phys. Rev. Lett.* **85**, 441 (2000).
 - [6] D. Gottesman and J. Preskill, *Phys. Rev. A* **63**, 022309 (2001).
 - [7] M. Curty, M. Lewenstein, and N. Lütkenhaus, *Phys. Rev. Lett.* **92**, 217903 (2003); see also quant-ph/0409047.
 - [8] A. Acin and N. Gisin, quant-ph/0310054.
 - [9] A. Acin, N. Gisin, and V. Scarani, *Quant. Info. Comp.* **3**, 563 (2003).
 - [10] H. F. Chau, *Phys. Rev. A* **66**, 060302 (2002).
 - [11] D. Gottesman and H.-K. Lo, *IEEE Trans. Inf. Theory* **49**, 457 (2003).
 - [12] C. H. Bennett, F. Bessette, G. Brassard, L. Salvail, and J. Smolin, *J. Cryptology* **5**, 3 (1992); A. Ekert and B. Huttner, *J. Mod. Opt.* **41**, 2455 (1994).
 - [13] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, *Rev. Mod. Phys.* **74**, 145 (2002).

- [14] D. Deutsch, A. Ekert, R. Jozsa, C. Macchiavello, S. Popescu, and A. Sanpera, Phys. Rev. Lett. **77**, 2818 (1996).
- [15] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A **54**, 3824 (1996); C. H. Bennett, Gilles Brassard, S. Popescu, B. Schumacher, J. A. Smolin and W. K. Wootters, Phys. Rev. Lett. **76**, 722 (1996).
- [16] The Bell states, $|\Phi^\pm\rangle \equiv \frac{1}{\sqrt{2}}(|0_A0_B\rangle \pm |1_A1_B\rangle)$ and $|\Psi^\pm\rangle \equiv \frac{1}{\sqrt{2}}(|0_A1_B\rangle \pm |1_A0_B\rangle)$, form an orthonormal basis in the two-qubit Hilbert space.
- [17] It is worth noting that a logarithmic scaling of the size of the random sample with the length of Alice's and Bob's key, is sufficient for security issues. See Ref. 18 for a rigorous proof. Nevertheless, throughout this work we will consider that half of the pairs are sacrificed for the verification test.
- [18] H.-K. Lo, H. F. Chau and M. Ardehali, quant-ph/0011056.
- [19] Note that in the absence of noise and eavesdropping each pair of qubits shared between Alice and Bob is in the Bell state $|\Phi^+\rangle$. Thus, in this ideal system, Alice and Bob obtain perfectly correlated measurement results whenever they perform their measurements along the same basis.
- [20] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996); M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 8 (1996).
- [21] N. Gisin and S. Wolf, Phys. Rev. Lett. **83**, 4200 (1999). See also quant-ph/0005042.
- [22] A. Acin, L. Masanes, and N. Gisin, Phys. Rev. Lett. **91**, 167901 (2003).